

On a theorem of Artin

Arjeh M. Cohen & Luis Paris

April 2002

Abstract

We determine the epimorphisms $A \rightarrow W$ from the Artin group A of type Γ onto the Coxeter group W of type Γ , in case Γ is an irreducible Coxeter graph of spherical type, and we prove that the kernel of the standard epimorphism is a characteristic subgroup of A . This generalizes an over 50 years old result of Artin.

AMS Subject Classification: Primary 20F36; Secondary 20F55. Keywords: Braid groups, Artin groups, Coxeter groups, characteristic subgroups, reflections.

1. Introduction

In 1947, Artin [Art] published the following two results on the braid group B_n on n strings.

Theorem 1.1 (Artin, [Art]). *Suppose that $\varphi : B_n \rightarrow \text{Sym}_n$ is an epimorphism. Then, up to an automorphism of Sym_n , the elements $\varphi(\sigma_i)$ ($i = 1, \dots, n-1$) are the standard generators $(i, i+1)$ of Sym_n , with the following two exceptions for $n = 4$:*

$$\varphi(\sigma_1) = (1, 2, 3, 4), \quad \varphi(\sigma_2) = (2, 1, 3, 4), \quad \varphi(\sigma_3) = (1, 2, 3, 4)$$

and

$$\varphi(\sigma_1) = (1, 2, 3, 4), \quad \varphi(\sigma_2) = (2, 1, 3, 4), \quad \varphi(\sigma_3) = (4, 3, 2, 1),$$

where σ_i denotes the i -th standard generator of B_n .

Theorem 1.2 (Artin, [Art]). *The pure braid group PB_n is a characteristic subgroup of B_n .*

Let Γ be a Coxeter graph with set of vertices $\{1, \dots, n\}$ (cf. [Bou] for terminology). For two objects a, b and for $m \in \mathbb{N}$, define the word

$$[a, b]^m = \begin{cases} (ab)^{\frac{m}{2}} & \text{if } m \text{ is even,} \\ (ab)^{\frac{m-1}{2}} a & \text{if } m \text{ is odd.} \end{cases}$$

The *Artin group* of type Γ is the group A generated by n elements $\sigma_1, \dots, \sigma_n$, and subject to the relations

$$[\sigma_i, \sigma_j]^{m_{i,j}} = [\sigma_j, \sigma_i]^{m_{i,j}}$$

for $1 \leq i < j \leq n$, and $m_{i,j} \neq \infty$, where $(m_{i,j})_{i,j}$ is the Coxeter matrix of Γ . The Coxeter system of type Γ is denoted by (W, S) with S consisting of the images s_i of σ_i ($i = 1, \dots, n$) under the natural epimorphism $\mu : A \rightarrow W$. The number of generators n is called the *rank* of the Artin group (and of the Coxeter group). We say that A is *irreducible* if Γ is connected, and that A is of *spherical type* if W is finite. Define the *colored Artin group* CA of type Γ to be the kernel of the standard epimorphism $\mu : A \rightarrow W$.

Call two epimorphisms $\varphi_1, \varphi_2 : A \rightarrow W$ *equivalent* if there is an automorphism α of W such that $\varphi_2 = \alpha \circ \varphi_1$. Members of the equivalence class of the standard epimorphism are called *ordinary* epimorphisms. The epimorphisms which are not ordinary are called *extraordinary*.

Note that the Artin group A of spherical type A_{n-1} is the braid group B_n of Theorem 1.1 and that the corresponding Coxeter group W coincides with Sym_n . In this case, CA is the pure braid group PB_{n-1} of Theorem 1.2. Thus, it makes sense to ask whether Artin's results for type A_{n-1} also hold for other Coxeter graphs. In this paper, we prove the results corresponding to Theorems 1.1 and 1.2 for the irreducible Artin groups of spherical type by proving the following two theorems.

Theorem 1.3. *Let A be an irreducible Artin group of spherical type distinct from A_3 with corresponding Coxeter group W . Then the equivalence classes of extraordinary epimorphisms $\varphi : A \rightarrow W$ are as given in Table 1.*

Coxeter type	epimorphism	defined in Proposition
$I_2(m)$ ($m \equiv 0 \pmod{4}$)	ν_m^1, ν_m^2	2.2
B_n (n even)	ν_n^1	3.2
B_n (n odd)	ν_n^1, ν_n^2	3.2
D_n (n odd)	ν_n^1, ν_n^2	4.2
H_3	ν_3^1	5.2

Table 1. The extraordinary epimorphisms up to equivalence.

Theorem 1.4. *Let A be an irreducible Artin group of spherical type. Then CA is a characteristic subgroup of A .*

Each ‘family’ of irreducible Artin groups of spherical type is treated separately. Our strategy for the proof of Theorem 1.3 is to analyze the epimorphisms $A \rightarrow W$ in much the same way Artin did for type A_n , and to study epimorphisms $A \rightarrow \text{Sym}_n$ for types B_n, D_n . In order to prove Theorem 1.4, we establish that the kernel of every surjective composition $\mu \circ \Phi$ of μ with an automorphism Φ of A coincides with CA . For this purpose, we study an action of A on the set $\mathbf{Z} \times T$, where T is the set of reflections of W , which is compatible (under μ) with the action of W on the set $\{\pm 1\} \times T$ of abstract walls of (W, S) (Lemma 3.7).

The groups of rank two are treated in Section 2, Artin groups of type B_n in Section 3, Artin groups of type D_n in Section 4, and Artin groups of type $H_3, H_4, F_4, E_6, E_7, E_8$ in Section 5. As noted above, type A_n is dealt with in [Art]. The classification of the epimorphisms $A \rightarrow W$ for A of type B_n and D_n is almost done in [Lin]; our contribution consists of Lemmas 3.5 and 4.4 and leads to a complete classification. Our approach to these two families is the same as Lin's, except that we do not use any homological argument.

Acknowledgments. The second author would like to thank Jean Michel who introduced him to GAP and to the package ‘Chevie’ of GAP.

2. Artin groups of rank two

We start this section with a lemma which will be used throughout the whole paper. Its proof is direct from the definition of a Coxeter group.

Lemma 2.1. *Let A be an Artin group of spherical type. If all the images of the standard generators under an epimorphism $\varphi : A \rightarrow W$ are involutions, then φ is ordinary. $\square$*

Recall from [Bou] that $I_2(m)$ denotes the Coxeter graph on two nodes which are joined by an m -valued edge. It is irreducible whenever $m \geq 3$. By abuse of notation, we will use $I_2(3), I_2(4)$, and $I_2(6)$ to denote the Coxeter graph of type A_2, B_2 , and G_2 , respectively.

Proposition 2.2. *Let $m \geq 3$. For the Coxeter graph $I_2(m)$, each epimorphism $A \rightarrow W$ is ordinary if $m \not\equiv 0 \pmod{4}$. If $m \equiv 0 \pmod{4}$, then there are two equivalence classes of extraordinary epimorphisms. These are represented by $\nu_m^1, \nu_m^2 : A \rightarrow W$, where*

$$\begin{aligned} \nu_m^1(\sigma_1) &= s_1, & \nu_m^1(\sigma_2) &= s_2 s_1, \\ \nu_m^2(\sigma_1) &= s_1 s_2, & \nu_m^2(\sigma_2) &= s_2. \end{aligned}$$

Proof. Let $r = s_2 s_1$. Let C be the subgroup of W generated by r . It is a cyclic normal subgroup of W of order m , and $W = C \rtimes \langle \sigma_1 \rangle \cong \mathbf{Z}/m\mathbf{Z} \rtimes \mathbf{Z}/2\mathbf{Z}$. Let $\varphi : A \rightarrow W$ be an epimorphism, and let $t_1 = \varphi(\sigma_1)$ and $t_2 = \varphi(\sigma_2)$. Four cases have to be considered.

Case 1: $t_1, t_2 \in s_1 C$. Then t_1 and t_2 are involutions, and φ is ordinary by Lemma 2.1.

Case 2: $t_1, t_2 \in C$. Then $\text{Im}(\varphi) \subset C$, and φ is not surjective. So, this case does not occur.

Case 3: $t_1 \in s_1 C$ and $t_2 \in C$. If m is odd, then t_1 and t_2 are conjugate (since σ_1 and σ_2 are conjugate), a contradiction with $t_2 \in C$ and $t_1 \notin C$ as C is a normal subgroup of W . So, m is even. The fact that t_1 and t_2 generate W implies that t_2 is of order m , so, up to an automorphism of W , we can assume that $t_1 = s_1$ and $t_2 = r$. Consider the equality

$$t_1 t_2 t_1 t_2 = s_1 r s_1 r = 1.$$

If $m \equiv 2 \pmod{4}$, then $[t_1, t_2]^m = [t_2, t_1]^m$ implies $s_1 r = r s_1$, but this last equality does not hold. Hence, $m \equiv 0 \pmod{4}$. Now φ is indeed a homomorphism as

$$[t_1, t_2]^m = [t_2, t_1]^m = 1,$$

and φ is equivalent to ν_m^1 .

Case 4: $t_1 \in C$ and $t_2 \in s_1 C$. Then Case 3 applies with the indices 1 and 2 interchanged. Therefore $m \equiv 0 \pmod{4}$, and φ is equivalent to ν_m^2 . $\square$

Recall that $\mu : A \rightarrow W$ denotes the standard epimorphism mapping σ_i to s_i .

Lemma 2.3. *If $\Phi \in \text{Aut}(A)$ is such that $\mu \circ \Phi$ is ordinary, then Φ preserves CA .*

Proof. As $\mu \circ \Phi : A \rightarrow W$ is ordinary, there is an automorphism α of W such that $\mu \circ \Phi = \alpha \circ \mu$. Now

$$\Phi^{-1}(CA) = \Phi^{-1}(\text{Ker}(\mu)) = \text{Ker}(\mu \circ \Phi) = \text{Ker}(\alpha \circ \mu) = \text{Ker}(\mu) = CA. \quad \square$$

Proposition 2.4. *Let $m \geq 3$. For the Coxeter graph $I_2(m)$, the colored Artin group CA is a characteristic subgroup of A .*

Proof. Let $\Phi : A \rightarrow A$ be an automorphism. In view of Lemma 2.3, we may assume that $\mu \circ \Phi$ is an extraordinary epimorphism. In particular, by Proposition 2.2, $m \equiv 0 \pmod{4}$. Let $\Delta = [\sigma_1, \sigma_2]^m \in A$, and let $\delta = [s_1, s_2]^m \in W$. Since m is even, by [BS], the center of A is the infinite cyclic subgroup generated by Δ , so $\Phi(\Delta) = \Delta^\varepsilon$, where $\varepsilon \in \{\pm 1\}$, whence

$$(\mu \circ \Phi)(\Delta) = \mu(\Delta^\varepsilon) = \delta^\varepsilon = \delta.$$

On the other hand, by Proposition 2.2, there exist an automorphism α of W and $i \in \{1, 2\}$ such that $\mu \circ \Phi = \alpha \circ \nu_m^i$. But


$$\nu_m^1(\Delta) = [s_1, s_2 s_1]^m = 1 \quad \text{and} \quad \nu_m^2(\Delta) = [s_1 s_2, s_2]^m = 1,$$

whence $\delta = \mu \circ \Phi(\Delta) = (\alpha \circ \nu_m^i)(\Delta) = 1$. This contradicts $\delta \neq 1$, so we are done. $\square$

3. Artin groups of type B_n

For a group G and an element $g \in G$, we denote by $\text{Conj}_g : G \rightarrow G$ the inner automorphism $h \mapsto ghg^{-1}$. Call two epimorphisms $\varphi_1, \varphi_2 : G \rightarrow H$ *conjugate* if there is $h \in H$ such that $\varphi_2 = \text{Conj}_h \circ \varphi_1$.

We shall label the nodes of the Coxeter graph B_n as follows:



Let A be an Artin group of type B_n , $n \geq 3$, and let W be its corresponding Coxeter group. The first step in our investigation of the epimorphisms $A \rightarrow W$ is to classify the epimorphisms $A \rightarrow \text{Sym}_n$ up to

conjugation. This has also been carried out by Zinde [Zin], and the answer is as follows. Define the standard epimorphism $\eta : A \rightarrow \text{Sym}_n$ by

$$\eta(\sigma_1) = 1, \quad \eta(\sigma_i) = (i-1, i) \quad \text{for } i = 2, \dots, n.$$

Proposition 3.1 (Zinde, [Zin]). *Let $n \geq 3$. For the Coxeter graph B_n , every epimorphism $A \rightarrow \text{Sym}_n$ is conjugate to η if $n \neq 4$ and $n \neq 6$.*

If $n = 4$, then there are precisely three conjugacy classes of epimorphisms; these are represented by $\eta, \zeta_4^1, \zeta_4^2 : A \rightarrow \text{Sym}_4$, where

$$\zeta_4^1(\sigma_1) = 1, \quad \zeta_4^1(\sigma_2) = (1, 2, 3, 4), \quad \zeta_4^1(\sigma_3) = (2, 1, 3, 4), \quad \zeta_4^1(\sigma_4) = (1, 2, 3, 4),$$

and

$$\zeta_4^2(\sigma_1) = 1, \quad \zeta_4^2(\sigma_2) = (1, 2, 3, 4), \quad \zeta_4^2(\sigma_3) = (2, 1, 3, 4), \quad \zeta_4^2(\sigma_4) = (4, 3, 2, 1).$$

If $n = 6$, then there are precisely two conjugacy classes of epimorphisms; these are represented by $\eta, \zeta_6 : A \rightarrow \text{Sym}_6$, where

$$\begin{aligned} \zeta_6(\sigma_1) &= 1, & \zeta_6(\sigma_2) &= (1, 2)(3, 4)(5, 6), & \zeta_6(\sigma_3) &= (1, 5)(2, 3)(4, 6), \\ \zeta_6(\sigma_4) &= (1, 3)(2, 4)(5, 6), & \zeta_6(\sigma_5) &= (1, 2)(3, 5)(4, 6), & \zeta_6(\sigma_6) &= (1, 4)(2, 3)(5, 6). \end{aligned}$$

Moreover, if we set

$$\begin{aligned} r_1 &= 1, & r_2 &= s_2 s_4 s_6, & r_3 &= s_2 s_3 s_2 s_5 s_4 s_3 s_2 s_6 s_5, \\ r_4 &= s_3 s_2 s_4 s_3 s_6, & r_5 &= s_2 s_5 s_4 s_6 s_5, & r_6 &= s_2 s_3 s_2 s_4 s_3 s_2 s_6, \end{aligned}$$

then $r_i = \zeta_6(\sigma_i)$ for all $i = 1, \dots, 6$, and the automorphism α of Sym_6 determined by $(i-1, i) \mapsto r_i$ ($i = 2, \dots, 6$) satisfies $\zeta_6 = \alpha \circ \eta$. $\square$

The proof of Proposition 3.1 can be recovered from Artin's proof of [Art, Thm.1] by an easy adaptation.

We need a more precise description of the well-known isomorphism $W \cong (\mathbf{Z}/2\mathbf{Z})^n \rtimes \text{Sym}_n$. Put $c_1 = s_1$ and, for $i = 2, \dots, n$, denote by c_i the element $s_i \cdots s_2 s_1 s_2 \cdots s_i$ of W . Then $C = \langle c_1, \dots, c_n \rangle \cong (\mathbf{Z}/2\mathbf{Z})^n$ is the kernel of the natural epimorphism $p : W = C \rtimes S \rightarrow \text{Sym}_n$, where $S = \langle s_2, \dots, s_n \rangle \cong \text{Sym}_n$. It is convenient to identify $c_1^{a_1} \cdots c_n^{a_n} \in C$ with $[a_1, \dots, a_n] \in (\mathbf{Z}/2\mathbf{Z})^n$, so that each element of W can be viewed as the product of an element from $(\mathbf{Z}/2\mathbf{Z})^n$ and a permutation in Sym_n . Observe that the longest element $\delta = (s_1 \cdots s_n)^n$ of W coincides with $c_1 \cdots c_n$.

Proposition 3.2. *Let $n \geq 3$. For the Coxeter graph B_n , representatives of the equivalence classes of extraordinary epimorphisms $A \rightarrow W$ are*

- (i) ν_n^1 , if n is even,
- (ii) ν_n^1 and ν_n^2 , if n is odd,

where ν_n^1 is the homomorphism $A \rightarrow W$ determined by

$$\begin{aligned} \nu_n^1(\sigma_1) &= c_1 \\ \nu_n^1(\sigma_i) &= c_{i-1} s_i \quad \text{for } i = 2, \dots, n, \end{aligned}$$

and, for n odd, ν_n^2 is the homomorphism $A \rightarrow W$ determined by

$$\begin{aligned} \nu_n^2(\sigma_1) &= \delta \\ \nu_n^2(\sigma_i) &= c_{i-1} s_i \quad \text{for } i = 2, \dots, n. \end{aligned}$$

Before proving Proposition 3.2, we establish three lemmas. Observe that $\eta = p \circ \mu$.

Lemma 3.3. *Let $\varphi : A \rightarrow W$ be an extraordinary epimorphism such that $p \circ \varphi$ is the standard epimorphism $\eta : A \rightarrow \text{Sym}_n$. Then, up to an automorphism of W , we have $\varphi(\sigma_i) = c_{i-1} \cdot s_i$ for $i = 2, \dots, n$.*

Proof. Let $t_i = \varphi(\sigma_i)$, $i = 1, 2, \dots, n$. Note that $t_1 \in C$ in view of the assumption that $p \circ \varphi = \eta$. Thus, either $t_1 = 1$ or t_1 is an involution. Note also that t_i is conjugate to t_2 for $i = 3, \dots, n$.

Assume first that t_2 is an involution. For $i = 3, \dots, n$, the element t_i , being conjugate to t_2 , is also an involution. Furthermore, t_1 cannot be the identity, for otherwise we would have an epimorphism $\text{Sym}_n \rightarrow W$, $s_i \mapsto t_{i+1}$ ($i = 1, \dots, n-1$). Hence, t_1 is also an involution, and, by Lemma 2.1, φ is ordinary. This is contrary to the hypotheses, so t_2 is not an involution.

As $p \circ \varphi = \eta$, for $i = 2, \dots, n$ we can write $t_i = v_i \cdot s_i$ with $v_i = c_1^{a_{i,1}} \cdots c_n^{a_{i,n}} \in C$. The inequalities $t_i^2 \neq 1$ imply $a_{i,i} \equiv 1 + a_{i,i-1} \pmod{2}$ for $i = 2, \dots, n$. The equalities $t_i t_2 = t_2 t_i$ imply $a_{2,i} \equiv a_{2,3} \pmod{2}$ for $i = 4, \dots, n$. Finally, the equalities $t_i t_{i+1} t_i = t_{i+1} t_i t_{i+1}$ ($i = 2, \dots, n-1$) imply $a_{i,j} \equiv a_{2,3} \pmod{2}$ for $i = 2, \dots, n$ and $j \neq i-1, i$. Write $a = a_{2,3}$ and $a_i = a_{i,i-1}$ for $i = 2, \dots, n$. Then

$$v_i = c_1^a \cdots c_{i-2}^a c_{i-1}^{a_i} c_i^{1+a_i} c_{i+1}^a \cdots c_n^a.$$

Suppose $a \equiv 1 \pmod{2}$. Let α_0 be the automorphism of W determined by

$$\begin{aligned} \alpha_0(s_1) &= s_1, \\ \alpha_0(s_i) &= \delta s_i \quad \text{for } i = 2, \dots, n. \end{aligned}$$

Then

$$(\alpha_0 \circ \varphi)(\sigma_i) = \alpha_0(t_i) = c_{i-1}^{1+a_i} c_i^{a_i} s_i \quad \text{for } i = 2, \dots, n,$$

so we can assume $a = 0$.

Suppose $a_n \equiv \cdots \equiv a_{i+1} \equiv 1 \pmod{2}$ and $a_i \equiv 0 \pmod{2}$. Then

$$\begin{aligned} (\text{Conj}_{c_{i-1}} \circ \varphi)(\sigma_j) &= \text{Conj}_{c_{i-1}}(t_j) = t_j \quad \text{if } j \neq i-1, i, \\ (\text{Conj}_{c_{i-1}} \circ \varphi)(\sigma_j) &= \text{Conj}_{c_{i-1}}(t_j) = c_{j-1} c_j t_j \quad \text{if } j = i-1, i, \end{aligned}$$

so we can assume $a_n \equiv \cdots \equiv a_{i+1} \equiv a_i \equiv 1 \pmod{2}$. An iteration of this argument shows that we can assume $a_i \equiv 1 \pmod{2}$ for $i = 2, \dots, n$. But then $v_i = c_{i-1}$. $\square$

Lemma 3.4. Let $\varphi : A \rightarrow W$ be an extraordinary epimorphism such that $p \circ \varphi = \eta$.

- (i) If n is even, then φ is equivalent to ν_n^1 .
- (ii) If n is odd, then φ is equivalent to ν_n^1 or to ν_n^2 .

Proof. Let $t_i = \varphi(\sigma_i)$, for $i = 1, 2, \dots, n$. By Lemma 3.3, we can assume $t_i = c_{i-1} \cdot s_i$ for $i = 2, \dots, n$. Write $t_1 = c_1^{a_1} c_2^{a_2} \cdots c_n^{a_n} \in C$. The equalities $t_i t_1 = t_1 t_i$ ($i = 3, \dots, n$) imply $a_i \equiv a_2 \pmod{2}$ for $i = 3, \dots, n$, so, $t_1 = c_1^{a_1} c_2^{a_2} \cdots c_n^{a_2}$.

Consider the epimorphism $\text{sg} : W \rightarrow \{\pm 1\}$, $s_i \mapsto -1$. It satisfies $\text{sg}(t_1) \equiv (-1)^{a_1 + (n-1)a_2}$ and $\text{sg}(t_i) = 1$ for $i = 2, \dots, n$. So, if $a_1 + (n-1)a_2 \equiv 0 \pmod{2}$, then $\text{Im}(\varphi) \subset \text{Ker}(\text{sg})$, and φ is not surjective. Therefore, $a_1 + (n-1)a_2 \equiv 1 \pmod{2}$. If $a_1 \equiv 1 \pmod{2}$ and $a_2 \equiv 0 \pmod{2}$, then $\varphi = \nu_n^1$, and if $a_1 \equiv 1 \pmod{2}$ and $a_2 \equiv 1 \pmod{2}$, then n is odd and $\varphi = \nu_n^2$. Assume $a_1 \equiv 0 \pmod{2}$ and $a_2 \equiv 1 \pmod{2}$. Then n is even. Let α_1 be the automorphism of W determined by

$$\begin{aligned} \alpha_1(s_1) &= \delta s_1, \\ \alpha_1(s_i) &= s_i \quad \text{for } i = 2, \dots, n. \end{aligned}$$

Then

$$\begin{aligned} (\alpha_1 \circ \varphi)(\sigma_1) &= \alpha_1(t_1) = c_1, \\ (\alpha_1 \circ \varphi)(\sigma_i) &= \alpha_1(t_i) = c_{i-1} \cdot s_i \quad \text{for } i = 2, \dots, n, \end{aligned}$$

so $\alpha_1 \circ \varphi = \nu_n^1$. $\square$

Lemma 3.5. For the Coxeter graph B_6 , there is no extraordinary epimorphism $\varphi : A \rightarrow W$ such that $p \circ \varphi = \zeta_6$.

For the Coxeter graph B_4 , there is no extraordinary epimorphism $\varphi : A \rightarrow W$ such that either $p \circ \varphi = \zeta_4^1$ or $p \circ \varphi = \zeta_4^2$.

Proof. The proof is a direct calculation. It can be carried out in GAP or Magma by means of a Todd Coxeter enumeration leading to a permutation representation of W , or by use of the package ‘Chevie’ of GAP. Here, we explain how we deal with ζ_6 . The other two cases can be handled similarly.

We first compute the set $X_i = \{xr_i; x \in C \text{ and } \text{order}(xr_i) \neq 2\}$, for $i = 2, \dots, 6$, where r_i is as in Proposition 3.1. We have $|X_i| = 56$ for each i . For $i = 3, \dots, 6$, let Y_i be the set of $(i-1)$ -tuples $(t_2, \dots, t_i)$ which satisfy

- $t_j \in X_j$, for $j = 2, \dots, i$,
- $t_j t_{j+1} t_j = t_{j+1} t_j t_{j+1}$, for $j = 2, \dots, i-1$,
- $t_j t_k = t_k t_j$, for $|j-k| \geq 2$.

We compute successively Y_3, Y_4, Y_5 , and Y_6 . We have $|Y_3| = 224$, $|Y_4| = 192$, $|Y_5| = 64$, and $Y_6 = \emptyset$. The last equality together with Lemma 2.1 proves the lemma. $\square$

Proof of Proposition 3.2. Let $\varphi : A \rightarrow W$ be an extraordinary epimorphism. By Proposition 3.1 and the fact that $p(W) = \text{Sym}_n$, we can assume that either $p \circ \varphi$ is the standard epimorphism, or $n = 6$ and $p \circ \varphi = \zeta_6$, or $n = 4$ and $p \circ \varphi = \zeta_4^i$, $i = 1, 2$. By Lemma 3.5, the last two cases do not occur, so $p \circ \varphi$ is standard. Now, Proposition 3.2 follows from Lemma 3.4. $\square$

Let A be an Artin group. By Lemma 2.3, in order to prove that CA is a characteristic subgroup of A , it suffices to show that, for a given automorphism $\Phi : A \rightarrow A$, the composition $\mu \circ \Phi : A \rightarrow W$ with the standard epimorphism $\mu : A \rightarrow W$ is ordinary. For example, for an Artin group of type B_n , we would like to prove that $\mu \circ \Phi \neq \nu_n^i$, for $i = 1, 2$. Our general strategy is to use the homomorphism $U : A \rightarrow \text{Sym}(\mathbf{Z} \times T)$ defined in Lemma 3.6 below.

Let $T = \{ws w^{-1}; w \in W \text{ and } s \in S\}$ be the set of reflections in W . For all $i = 1, \dots, n$, define the transformation $u_i : \{\pm 1\} \times T \rightarrow \{\pm 1\} \times T$ by

$$u_i(\varepsilon, t) = \begin{cases} (\varepsilon, s_i t s_i) & \text{if } t \neq s_i, \\ (-\varepsilon, t) & \text{if } t = s_i. \end{cases}$$

The following lemma can be found in [Bou, Ch.4, $n^\circ 1.4$].

Lemma 3.6 (Bourbaki, [Bou]). *For each type Γ , the assignment $s_i \mapsto u_i$ determines a homomorphism $u : W \rightarrow \text{Sym}(\{\pm 1\} \times T)$, where $\text{Sym}(\{\pm 1\} \times T)$ denotes the group of permutations of $\{\pm 1\} \times T$. $\square$*

Now, for $i = 1, \dots, n$, define $U_i : \mathbf{Z} \times T \rightarrow \mathbf{Z} \times T$ by

$$U_i(k, t) = \begin{cases} (k, s_i t s_i) & \text{if } t \neq s_i \\ (k+1, t) & \text{if } t = s_i \end{cases}$$

Observe that each U_i is a permutation of $\mathbf{Z} \times T$ with inverse given by the same formula as for U_i , but with $k+1$ replaced by $k-1$. The lemma below is easily derived by arguments similar to those of [Bou, Ch.4, $n^\circ 1.4$].

Lemma 3.7. *For each type Γ , the permutations U_i of $\mathbf{Z} \times T$ satisfy the following properties.*

- (1) *The assignment $\sigma_i \mapsto U_i$ determines a unique homomorphism $U : A \rightarrow \text{Sym}(\mathbf{Z} \times T)$.*
- (2) *Let $g \in A$, and let $(k, r) \in \mathbf{Z} \times T$. Write $U(g)(k, r) = (k', r')$. Then*

$$u(\mu(g))((-1)^k, r) = ((-1)^{k'}, r').$$

- (3) *For $k \in \mathbf{Z}$ the element $L_k \in \text{Sym}(\mathbf{Z} \times T)$ defined by $L_k(l, t) = (k+l, t)$ commutes with each U_i . In particular, for $g \in A$ we have $U(g)(k, t) = L_k U(g)(0, t)$. $\square$*

A direct consequence of the second part of this lemma is:

Corollary 3.8. *Let $g_1, g_2 \in A$, and let $(k, r) \in \mathbf{Z} \times T$. Write $(k_1, r_1) = U(g_1)(k, r)$, and $(k_2, r_2) = U(g_2)(k, r)$. If $\mu(g_1) = \mu(g_2)$, then $r_1 = r_2$ and $k_1 \equiv k_2 \pmod{2}$. $\square$*

We now prove the second main result for B_n .

Proposition 3.9. *Let $n \geq 3$. For the Coxeter graph B_n , the colored Artin group CA is a characteristic subgroup of A .*

The key of the proof of Proposition 3.9 is the following lemma.

Lemma 3.10. *Let A be an Artin group of type B_n , $n \geq 3$, and let $\Phi : A \rightarrow A$ be a homomorphism. Then $\mu \circ \Phi \neq \nu_n^1$, and $\mu \circ \Phi \neq \nu_n^2$ (if n is odd).*

Proof. Assume that there exists a homomorphism $\Phi : A \rightarrow A$ such that $\mu \circ \Phi = \nu_n^i$, where $i \in \{1, 2\}$. Consider the following elements of T .

$$\begin{aligned} r_1 &= s_2, & r_2 &= s_2 s_3 s_2, & r_3 &= s_3, \\ r_4 &= s_1 s_2 s_1, & r_5 &= s_1 s_2 s_3 s_2 s_1, & r_6 &= s_2 s_1 s_2 s_3 s_2 s_1 s_2. \end{aligned}$$

Put

$$\tau_2 = \sigma_1 \sigma_2, \quad \text{and} \quad \tau_3 = \sigma_2 \sigma_1 \sigma_2 \sigma_3,$$

so that $\mu(\tau_j) = \nu_n^i(\sigma_j)$ for $j = 2, 3$. Straightforward calculations give

$$\begin{aligned} U(\tau_2)(0, r_1) &= (1, r_4), & U(\tau_2)(0, r_2) &= (0, r_3), & U(\tau_2)(0, r_3) &= (0, r_5), \\ U(\tau_2)(0, r_4) &= (0, r_1), & U(\tau_2)(0, r_5) &= (0, r_6), & U(\tau_2)(0, r_6) &= (0, r_2). \end{aligned}$$

and

$$\begin{aligned} U(\tau_3)(0, r_1) &= (0, r_2), & U(\tau_3)(0, r_2) &= (1, r_4), & U(\tau_3)(0, r_3) &= (1, r_6), \\ U(\tau_3)(0, r_4) &= (0, r_5), & U(\tau_3)(0, r_5) &= (1, r_1), & U(\tau_3)(0, r_6) &= (0, r_3). \end{aligned}$$

Now $\mu(\tau_2) = \nu_n^i(\sigma_2) = \mu(\Phi(\sigma_2))$ and $\mu(\tau_3) = \nu_n^i(\sigma_3) = \mu(\Phi(\sigma_3))$, so, by Corollary 3.8, there exist integers $a_1, \dots, a_6, b_1, \dots, b_6$ such that

$$\begin{aligned} U(\Phi(\sigma_2))(0, r_1) &= (2a_1 + 1, r_4), & U(\Phi(\sigma_2))(0, r_2) &= (2a_2, r_3), & U(\Phi(\sigma_2))(0, r_3) &= (2a_3, r_5), \\ U(\Phi(\sigma_2))(0, r_4) &= (2a_4, r_1), & U(\Phi(\sigma_2))(0, r_5) &= (2a_5, r_6), & U(\Phi(\sigma_2))(0, r_6) &= (2a_6, r_2). \end{aligned}$$

and

$$\begin{aligned} U(\Phi(\sigma_3))(0, r_1) &= (2b_1, r_2), & U(\Phi(\sigma_3))(0, r_2) &= (2b_2 + 1, r_4), & U(\Phi(\sigma_3))(0, r_3) &= (2b_3 + 1, r_6), \\ U(\Phi(\sigma_3))(0, r_4) &= (2b_4, r_5), & U(\Phi(\sigma_3))(0, r_5) &= (2b_5 + 1, r_1), & U(\Phi(\sigma_3))(0, r_6) &= (2b_6, r_3). \end{aligned}$$

Using Lemma 3.7(3), we derive

$$\begin{aligned} U(\Phi(\sigma_2 \sigma_3 \sigma_2))(0, r_1) &= (2a_1 + 2a_5 + 2b_4 + 1, r_6), & U(\Phi(\sigma_2 \sigma_3 \sigma_2))(0, r_2) &= (2a_2 + 2a_6 + 2b_3 + 1, r_2), \\ U(\Phi(\sigma_2 \sigma_3 \sigma_2))(0, r_3) &= (2a_1 + 2a_3 + 2b_5 + 2, r_4), & U(\Phi(\sigma_2 \sigma_3 \sigma_2))(0, r_4) &= (2a_2 + 2a_4 + 2b_1, r_3), \\ U(\Phi(\sigma_2 \sigma_3 \sigma_2))(0, r_5) &= (2a_3 + 2a_5 + 2b_6, r_5), & U(\Phi(\sigma_2 \sigma_3 \sigma_2))(0, r_6) &= (2a_4 + 2a_6 + 2b_2 + 1, r_1). \end{aligned}$$

and

$$\begin{aligned} U(\Phi(\sigma_3 \sigma_2 \sigma_3))(0, r_1) &= (2a_2 + 2b_1 + 2b_3 + 1, r_6), & U(\Phi(\sigma_3 \sigma_2 \sigma_3))(0, r_2) &= (2a_4 + 2b_1 + 2b_2 + 1, r_2), \\ U(\Phi(\sigma_3 \sigma_2 \sigma_3))(0, r_3) &= (2a_6 + 2b_2 + 2b_3 + 2, r_4), & U(\Phi(\sigma_3 \sigma_2 \sigma_3))(0, r_4) &= (2a_5 + 2b_4 + 2b_6, r_3), \\ U(\Phi(\sigma_3 \sigma_2 \sigma_3))(0, r_5) &= (2a_1 + 2b_4 + 2b_5 + 2, r_5), & U(\Phi(\sigma_3 \sigma_2 \sigma_3))(0, r_6) &= (2a_3 + 2b_5 + 2b_6 + 1, r_1). \end{aligned}$$

The equalities $U(\Phi(\sigma_2 \sigma_3 \sigma_2))(0, r_i) = U(\Phi(\sigma_3 \sigma_2 \sigma_3))(0, r_i)$ for $i = 2, 3, 4, 5$ are equivalent to the equations

$$\begin{aligned} a_2 - a_4 + a_6 - b_1 - b_2 + b_3 &= 0, \\ a_1 + a_3 - a_6 - b_2 - b_3 + b_5 &= 0, \\ a_2 + a_4 - a_5 + b_1 - b_4 - b_6 &= 0, \\ -a_1 + a_3 + a_5 - b_4 - b_5 + b_6 &= 1. \end{aligned}$$

The sum of these four equations gives $2(a_2 + a_3 - b_2 - b_4) = 1$ which clearly is impossible. $\square$

Proof of Proposition 3.9. Let $\Delta = (\sigma_1 \cdots \sigma_n)^n \in A$, and recall that $\delta = (s_1 \cdots s_n)^n \in W$. Let $\alpha_0 : W \rightarrow W$ be the automorphism determined by

$$\alpha_0(s_1) = s_1, \quad \alpha_0(s_i) = \delta s_i \quad \text{for } i = 2, \dots, n,$$

and let $\Psi_0 : A \rightarrow A$ be the homomorphism determined by

$$\Psi_0(\sigma_1) = \sigma_1, \quad \Psi_0(\sigma_i) = \Delta \sigma_i \quad \text{for } i = 2, \dots, n.$$

Then $\mu \circ \Psi_0 = \alpha_0 \circ \mu$. For n even, let $\alpha_1 : W \rightarrow W$ be the automorphism determined by

$$\alpha_1(s_1) = \delta s_1, \quad \alpha_1(s_i) = s_i \quad \text{for } i = 2, \dots, n,$$

and let $\Psi_1 : A \rightarrow A$ be the homomorphism determined by


$$\Psi_1(\sigma_1) = \Delta \sigma_1, \quad \Psi_1(\sigma_i) = \sigma_i \quad \text{for } i = 2, \dots, n.$$

Then $\mu \circ \Psi_1 = \alpha_1 \circ \mu$. Observe also that, if $w \in W$, then $\text{Conj}_w \circ \mu = \mu \circ \text{Conj}_g$, where $g \in A$ is such that $\mu(g) = w$. By [Fra], $\text{Aut}(W)$ is generated as a monoid by $\{\text{Conj}_w; w \in W\} \cup \{\alpha_0, \alpha_1\}$ if n is even, and by $\{\text{Conj}_w; w \in W\} \cup \{\alpha_0\}$ if n is odd. In particular, by the above observations, if α is an automorphism of W , then there exists a homomorphism $\Psi : A \rightarrow A$ such that $\mu \circ \Psi = \alpha \circ \mu$.

Now, let $\Phi : A \rightarrow A$ be an automorphism. In order to prove Proposition 3.9, it suffices to show that $\mu \circ \Phi$ cannot be extraordinary. Suppose that $\mu \circ \Phi$ is extraordinary. By Proposition 3.2, there exist an automorphism $\alpha : W \rightarrow W$ and $i \in \{1, 2\}$ such that $\mu \circ \Phi = \alpha \circ \nu_n^i$. By the preceding argument, one can find a homomorphism $\Psi : A \rightarrow A$ such that $\mu \circ \Psi = \alpha^{-1} \circ \mu$, so $\mu \circ (\Psi \circ \Phi) = \nu_n^i$. But this equality contradicts Lemma 3.10. $\square$

4. Artin groups of type D_n

The strategy for classifying the epimorphisms $A \rightarrow W$ for the Artin groups of type D_n , $n \geq 4$, is the same as the strategy for the Artin groups of type B_n . First, we classify the epimorphisms $A \rightarrow \text{Sym}_n$ up to conjugation. This has also been carried out by Zinde [Zin], and the answer is as follows. Label the nodes of the Coxeter graph D_n as follows:



Define the standard epimorphism $\eta : A \rightarrow \text{Sym}_n$ as the homomorphism determined by

$$\begin{aligned} \eta(\sigma_1) &= \eta(\sigma_2) = (1, 2), \\ \eta(\sigma_i) &= (i-1, i) \quad \text{for } i = 3, \dots, n. \end{aligned}$$

Proposition 4.1 (Zinde, [Zin]). *Let $n \geq 4$. For the Coxeter graph D_n , every epimorphism $A \rightarrow \text{Sym}_n$ is conjugate to η if $n \neq 4$ and $n \neq 6$.*

If $n = 6$, then there are precisely two conjugacy classes of epimorphisms. These are represented by $\eta, \zeta_6 : A \rightarrow \text{Sym}_6$, respectively, where

$$\begin{aligned} \zeta_6(\sigma_1) &= (1, 2)(3, 4)(5, 6), & \zeta_6(\sigma_2) &= (1, 2)(3, 4)(5, 6), & \zeta_6(\sigma_3) &= (2, 3)(1, 5)(4, 6), \\ \zeta_6(\sigma_4) &= (3, 1)(2, 4)(5, 6), & \zeta_6(\sigma_5) &= (1, 2)(3, 5)(4, 6), & \zeta_6(\sigma_6) &= (2, 3)(1, 4)(5, 6). \end{aligned}$$

If $n = 4$, for each epimorphism $\varphi : A \rightarrow \text{Sym}_4$, there is a diagram automorphism Υ of A such that $\varphi \circ \Upsilon$ is conjugate to one of the three epimorphisms $\eta, \zeta_4^1, \zeta_4^2 : A \rightarrow \text{Sym}_4$, where

$$\zeta_4^1(\sigma_1) = \zeta_4^1(\sigma_2) = \zeta_4^1(\sigma_4) = (1, 2, 3, 4), \quad \zeta_4^1(\sigma_3) = (2, 1, 3, 4),$$

and

$$\zeta_4^2(\sigma_1) = \zeta_4^2(\sigma_2) = (1, 2, 3, 4), \quad \zeta_4^2(\sigma_3) = (2, 1, 3, 4), \quad \zeta_4^2(\sigma_4) = (4, 3, 2, 1). \quad \square$$

It is well known that W embeds into the Weyl group of type B_n with image an index two subgroup. The embedding can be made explicit by means of the isomorphism of the Weyl group of type B_n with $(\mathbf{Z}/2\mathbf{Z})^n \rtimes \text{Sym}_n$ described in the discussion preceding Proposition 3.2. The embedding maps s_1 to $(1, 1, 0, \dots, 0) \cdot (1, 2)$ and maps s_i to $(i-1, i)$ for $i \geq 2$. So, if we denote by H the index 2 subgroup of $(\mathbf{Z}/2\mathbf{Z})^n$ of all elements $(a_1, \dots, a_n)$ with $a_1 + \dots + a_n = 0$, we find that the image is $H \rtimes \text{Sym}_n$. Let $\delta = (s_1 s_2)(s_3 s_1 s_2 s_3) \cdots (s_n \cdots s_3 s_1 s_2 s_3 \cdots s_n)$ denote the longest element of W . Then δ is mapped to $(1, 1, \dots, 1) \in H$ if n is even and to $(0, 1, \dots, 1) \in H$ if n is odd. For n odd, write $u_1 = \delta$ and $u_i = s_i s_{i-1} \cdots s_2 u_1 s_2 \cdots s_{i-1} s_i$ if $i > 1$. We use Proposition 4.1 to prove Theorem 1.3 for D_n .

Proposition 4.2. *Let $n \geq 4$. For the Coxeter graph D_n , there are no extraordinary epimorphisms $A \rightarrow W$ if n is even. If n is odd, there are precisely two equivalence classes of extraordinary epimorphisms $A \rightarrow W$. These are represented by $\nu_n^1, \nu_n^2 : A \rightarrow W$, where*

$$\begin{aligned} \nu_n^1(\sigma_1) &= \nu_n^1(\sigma_2) = s_2 \delta = u_2 s_2, \\ \nu_n^1(\sigma_i) &= u_i s_i \quad \text{for } i = 2, \dots, n. \end{aligned}$$

and

$$\begin{aligned} \nu_n^2(\sigma_1) &= u_1 s_2, \\ \nu_n^2(\sigma_i) &= u_i s_i \quad \text{for } i = 2, \dots, n. \end{aligned}$$

The following two lemmas are preparations to the proof of Proposition 4.2. Denote by $p : W \rightarrow \text{Sym}_n$ the natural projection with kernel H .

Lemma 4.3. *Let $\varphi : A \rightarrow W$ be an extraordinary epimorphism such that $p \circ \varphi$ is the standard epimorphism $\eta : A \rightarrow \text{Sym}_n$. Then n is odd and φ is equivalent to ν_n^1 or ν_n^2 .*

Proof. Let $t_i = \varphi(\sigma_i)$ for $i = 1, 2, \dots, n$. Write $t_1 = v_1 \cdot (1, 2)$, and $t_i = v_i \cdot (i-1, i)$ (for $i = 2, \dots, n$), where $v_i \in H$. The same argument as in the proof of Lemma 3.3 shows that there exist $a, a_1, a_2, \dots, a_n \in \mathbf{Z}/2\mathbf{Z}$ such that $v_1 = (a_1, 1 + a_1, a, \dots, a)$, and $v_i = (a, \dots, a, a_i, 1 + a_i, a, \dots, a)$ (a_i being in the $(i-1)$ -st entry) for $i = 2, \dots, n$. The condition $v_i \in H$ implies that n is odd and $a = 1$. Using again the same argument as in the proof of Lemma 3.3, we see that $a_i = 1$ for all $i = 2, \dots, n$ (not necessarily for $i = 1$), up to an automorphism of W . $\square$

Lemma 4.4. *For the Coxeter graph D_6 , there is no extraordinary epimorphism $\varphi : A \rightarrow W$ such that $p \circ \varphi = \zeta_6$.*

For the Coxeter graph D_4 , there is no extraordinary epimorphism $\varphi : A \rightarrow W$ such that either $p \circ \varphi = \zeta_4^1$ or $p \circ \varphi = \zeta_4^2$.

Proof. The proof is a straightforward calculation which can be carried out in GAP (for instance with the package ‘Chevie’ of GAP) or Magma. It follows the same scheme as the proof of Lemma 3.5. $\square$

Proof of Proposition 4.2. Let $\varphi : A \rightarrow W$ be an extraordinary epimorphism. By Proposition 4.1, we can assume that either $p \circ \varphi$ is the standard epimorphism, or $n = 6$ and $p \circ \varphi = \zeta_6$, or $n = 4$ and $p \circ \varphi \circ \Upsilon = \zeta_4^i$, where Υ is a diagram automorphism of A , and $i \in \{1, 2\}$. By Lemma 4.4, the last two cases cannot happen, so $p \circ \varphi$ is standard. Proposition 4.2 now follows from Lemma 4.3. $\square$

We turn now to the proof of the second main result for D_n .

Proposition 4.5. *Let $n \geq 4$. For the Coxeter graph D_n , the colored Artin group CA is a characteristic subgroup of A .*

Remark. By Artin's Theorem (1.1), for the Coxeter graph A_n , each epimorphism $A \rightarrow W$ is standard, except for $n = 3$. In particular, this shows that, for an Artin group of type A_n , $n \neq 3$, the colored Artin group CA is a characteristic subgroup of A . Now, the proof given below also applies to A_3 , viewed as the Coxeter graph D_3 , and so provides an alternative proof to Theorem 1.2 for $n = 4$.

The following lemma is a preparation to the proof of Proposition 4.5.

Lemma 4.6. *Assume n is odd. Let A be an Artin group of type D_n , and let $\Phi : A \rightarrow A$ be a homomorphism. Then $\mu \circ \Phi \neq \nu_n^1$, and $\mu \circ \Phi \neq \nu_n^2$.*

Proof. Assume that there exists a homomorphism $\Phi : A \rightarrow A$ such that $\mu \circ \Phi = \nu_n^i$, where $i \in \{1, 2\}$. Consider the following six reflections.

$$\begin{aligned} r_1 &= s_1, & r_2 &= s_3 s_1 s_3, & r_3 &= s_2 s_3 s_1 s_3 s_2, \\ r_4 &= s_3, & r_5 &= s_2 s_3 s_2, & r_6 &= s_2. \end{aligned}$$

Let

$$\Delta = (\sigma_1 \sigma_2)(\sigma_3 \sigma_1 \sigma_2 \sigma_3) \cdots (\sigma_n \cdots \sigma_3 \sigma_1 \sigma_2 \sigma_3 \cdots \sigma_n).$$

Recall the map U defined in Lemma 3.7. The fact that $\delta = (s_1 s_2)(s_3 s_1 s_2 s_3) \cdots (s_n \cdots s_3 s_1 s_2 s_3 \cdots s_n)$ is a reduced expression for the longest element of W implies that $U(\Delta)(0, r_k) = (1, \delta r_k \delta)$ for $k = 1, \dots, 6$. Furthermore, since n is odd, $\delta s_1 \delta = s_2$, $\delta s_2 \delta = s_1$, and $\delta s_j \delta = s_j$ for $j = 3, \dots, n$. Let

$$\tau_2 = \sigma_2 \Delta, \quad \tau_3 = \sigma_3 \sigma_2 \Delta \sigma_2,$$

so that $\mu(\tau_2) = s_2 \delta = \nu_n^i(\sigma_2)$ and $\mu(\tau_3) = s_3 s_2 \delta s_2 = \nu_n^i(\sigma_3)$. A straightforward calculation gives

$$\begin{aligned} U(\tau_2)(0, r_1) &= (2, r_6), & U(\tau_2)(0, r_2) &= (1, r_4), & U(\tau_2)(0, r_3) &= (1, r_2), \\ U(\tau_2)(0, r_4) &= (1, r_5), & U(\tau_2)(0, r_5) &= (1, r_3), & U(\tau_2)(0, r_6) &= (1, r_1). \end{aligned}$$

and

$$\begin{aligned} U(\tau_3)(0, r_1) &= (2, r_5), & U(\tau_3)(0, r_2) &= (1, r_1), & U(\tau_3)(0, r_3) &= (2, r_4), \\ U(\tau_3)(0, r_4) &= (1, r_3), & U(\tau_3)(0, r_5) &= (1, r_6), & U(\tau_3)(0, r_6) &= (2, r_2). \end{aligned}$$

Now $\mu(\tau_2) = \nu_n^i(\sigma_2) = \mu(\Phi(\sigma_2))$ and $\mu(\tau_3) = \nu_n^i(\sigma_3) = \mu(\Phi(\sigma_3))$ so, by Corollary 3.8, there exist integers $a_1, \dots, a_6, b_1, \dots, b_6$ such that

$$\begin{aligned} U(\Phi(\sigma_2))(0, r_1) &= (2a_1, r_6), & U(\Phi(\sigma_2))(0, r_2) &= (2a_2 + 1, r_4), & U(\Phi(\sigma_2))(0, r_3) &= (2a_3 + 1, r_2), \\ U(\Phi(\sigma_2))(0, r_4) &= (2a_4 + 1, r_5), & U(\Phi(\sigma_2))(0, r_5) &= (2a_5 + 1, r_3), & U(\Phi(\sigma_2))(0, r_6) &= (2a_6 + 1, r_1). \end{aligned}$$

and

$$\begin{aligned} U(\Phi(\sigma_3))(0, r_1) &= (2b_1, r_5), & U(\Phi(\sigma_3))(0, r_2) &= (2b_2 + 1, r_1), & U(\Phi(\sigma_3))(0, r_3) &= (2b_3, r_4), \\ U(\Phi(\sigma_3))(0, r_4) &= (2b_4 + 1, r_3), & U(\Phi(\sigma_3))(0, r_5) &= (2b_5 + 1, r_6), & U(\Phi(\sigma_3))(0, r_6) &= (2b_6, r_2). \end{aligned}$$

We find that

$$\begin{aligned} U(\Phi(\sigma_2 \sigma_3 \sigma_2))(0, r_1) &= (2a_1 + 2a_2 + 2b_6 + 1, r_4), & U(\Phi(\sigma_2 \sigma_3 \sigma_2))(0, r_2) &= (2a_2 + 2a_3 + 2b_4 + 3, r_2), \\ U(\Phi(\sigma_2 \sigma_3 \sigma_2))(0, r_3) &= (2a_1 + 2a_3 + 2b_2 + 2, r_6), & U(\Phi(\sigma_2 \sigma_3 \sigma_2))(0, r_4) &= (2a_4 + 2a_6 + 2b_5 + 3, r_1), \\ U(\Phi(\sigma_2 \sigma_3 \sigma_2))(0, r_5) &= (2a_4 + 2a_5 + 2b_3 + 2, r_5), & U(\Phi(\sigma_2 \sigma_3 \sigma_2))(0, r_6) &= (2a_5 + 2a_6 + 2b_1 + 2, r_3). \end{aligned}$$

and

$$\begin{aligned} U(\Phi(\sigma_3 \sigma_2 \sigma_3))(0, r_1) &= (2a_5 + 2b_1 + 2b_3 + 1, r_4), & U(\Phi(\sigma_3 \sigma_2 \sigma_3))(0, r_2) &= (2a_1 + 2b_2 + 2b_6 + 1, r_2), \\ U(\Phi(\sigma_3 \sigma_2 \sigma_3))(0, r_3) &= (2a_4 + 2b_3 + 2b_5 + 2, r_6), & U(\Phi(\sigma_3 \sigma_2 \sigma_3))(0, r_4) &= (2a_3 + 2b_2 + 2b_4 + 3, r_1), \\ U(\Phi(\sigma_3 \sigma_2 \sigma_3))(0, r_5) &= (2a_6 + 2b_1 + 2b_5 + 2, r_5), & U(\Phi(\sigma_3 \sigma_2 \sigma_3))(0, r_6) &= (2a_2 + 2b_4 + 2b_6 + 2, r_3). \end{aligned}$$

The equalities $U(\Phi(\sigma_2\sigma_3\sigma_2))(0, r_i) = U(\Phi(\sigma_3\sigma_2\sigma_3))(0, r_i)$ for $i = 1, 2, 4, 5$ are equivalent to the equations

$$\begin{aligned} a_1 + a_2 - a_5 - b_1 - b_3 + b_6 &= 0 \\ -a_1 + a_2 + a_3 - b_2 + b_4 - b_6 &= -1 \\ -a_3 + a_4 + a_6 - b_2 - b_4 + b_5 &= 0 \\ a_4 + a_5 - a_6 - b_1 + b_3 - b_5 &= 0 \end{aligned}$$

The sum of these four equations gives $2(a_2 + a_4 - b_1 - b_2) = -1$, a contradiction. $\square$

Proof of Proposition 4.5. Let $\Phi : A \rightarrow A$ be an automorphism. Suppose that $\mu \circ \Phi$ is extraordinary. By Proposition 4.2, n is odd, and there exist an automorphism $\alpha : W \rightarrow W$ and $i \in \{1, 2\}$ such that $\mu \circ \Phi = \alpha \circ \nu_n^i$. By [Fra], $\text{Aut}W = \{\text{Conj}_w; w \in W\}$. Let $w \in W$ be such that $\alpha^{-1} = \text{Conj}_w$, and take $g \in A$ such that $\mu(g) = w$. Then $\alpha^{-1} \circ \mu = \text{Conj}_w \circ \mu = \mu \circ \text{Conj}_g$, whence $\mu \circ (\text{Conj}_g \circ \Phi) = \nu_n^i$. This equality contradicts Lemma 4.6. Hence $\mu \circ \Phi$ is ordinary, and so the proposition follows from Lemma 2.3. $\square$

5. The remainder

In this section, we classify up to equivalence the epimorphisms $A \rightarrow W$, and we prove that CA is a characteristic subgroup of A , for the Artin groups of type $H_3, H_4, F_4, E_6, E_7, E_8$. This finishes our proofs of Theorems 1.3 and 1.4. We start with the Artin group of type H_3 , which requires a separate treatment, because it is the only Artin group among those of type $H_3, H_4, F_4, E_6, E_7, E_8$ which has extraordinary epimorphisms. For the corresponding Coxeter diagrams, we adopt the labelings

$$\begin{array}{c} \circ \quad 5 \quad \circ \\ | \quad \quad | \\ 1 \quad \quad 2 \quad \quad 3 \end{array} \quad \text{and} \quad \begin{array}{c} \circ \quad 5 \quad \circ \quad \circ \\ | \quad \quad | \quad \quad | \\ 1 \quad \quad 2 \quad \quad 3 \quad \quad 4 \end{array}$$

for H_3 and H_4 and the labeling of [Bou] for the remaining types.

Lemma 5.1. *For the Coxeter graph H_3 , there are precisely two conjugacy classes of extraordinary epimorphisms $A \rightarrow W$. These are represented by $\nu_3^1, \nu_3^2 : A \rightarrow W$, where*

$$\nu_3^1(\sigma_1) = s_1 s_2 s_3, \quad \nu_3^1(\sigma_2) = s_2 s_1 s_2 s_3 s_2, \quad \nu_3^1(\sigma_3) = s_3 s_2 s_1,$$

and

$$\nu_3^2(\sigma_1) = s_1 s_2 s_1 s_2 s_3 s_2 s_1 s_2 s_3, \quad \nu_3^2(\sigma_2) = s_2 s_1 s_2 s_3 s_2 s_1 s_2 s_1 s_3 s_2 s_1 s_2 s_3, \quad \nu_3^2(\sigma_3) = s_2 s_3 s_2 s_1 s_2 s_1 s_3 s_2 s_1.$$

Proof. The proof is a straightforward calculation, which can be made in GAP (for instance, with the package ‘Chevie’) or Magma. First, we calculate a representative for each conjugacy class of W . Let C_1 be the set of these representatives. We compute $C_2 = \{x \in C_1; \text{sg}(x) = -1 \text{ and } \text{order}(x) \neq 2\}$. We have $|C_2| = 3$. Let

$$\begin{aligned} X_1 &= \{(t_1, t_2) \in C_2 \times W; t_1 \neq t_2, t_1 t_2 t_1 t_2 t_1 = t_2 t_1 t_2 t_1 t_2\}, \\ X_2 &= \{(t_1, t_2, t_3) \in X_1 \times W; t_1 t_3 = t_3 t_1, t_2 t_3 t_2 = t_3 t_2 t_3\}. \end{aligned}$$

We successively compute X_1 and X_2 . We have $|X_1| = 16$, and $|X_2| = 10$. We verify that, if $(t_1, t_2, t_3) \in X_2$, then $\{t_1, t_2, t_3\}$ generates W . Let $\sim$ be the equivalence relation on X_2 defined by $(t_1, t_2, t_3) \sim (t'_1, t'_2, t'_3)$ if there exists $w \in W$ such that $t'_i = wt_i w^{-1}$ for all $i = 1, 2, 3$. We compute a representative for each equivalence class. Let Y be the set of these representatives. Then

$$\begin{aligned} Y &= \{(s_1 s_2 s_3, s_2 s_1 s_2 s_3 s_2, s_3 s_2 s_1), \\ &\quad (s_1 s_2 s_1 s_2 s_3 s_2 s_1 s_2 s_3, s_2 s_1 s_2 s_3 s_2 s_1 s_2 s_1 s_3 s_2 s_1 s_2 s_3, s_2 s_3 s_2 s_1 s_2 s_1 s_3 s_2 s_1)\}. \end{aligned} \quad \square$$

Proposition 5.2. *For the Coxeter graph H_3 , there is a unique equivalence class of extraordinary epimorphisms $A \rightarrow W$. It is represented by $\nu_3^1 : A \rightarrow W$.*

Proof. Let $\alpha : W \rightarrow W$ be the automorphism determined by

$$\alpha(s_1) = s_1 s_2 s_3 s_2 s_1, \quad \alpha(s_2) = s_2 s_1 s_2 s_3 s_2 s_1 s_2, \quad \alpha(s_3) = s_1 s_2 s_1 s_2 s_1.$$

Then $\alpha \circ \nu_3^1 = \nu_3^2$. $\square$

Proposition 5.3. *For the Coxeter graph H_3 , the colored Artin group CA is a characteristic subgroup of A .*

Proof. Let $\Phi : A \rightarrow A$ be an automorphism. In order to prove Proposition 5.3, it suffices to show that $\mu \circ \Phi$ is an ordinary epimorphism, cf. Lemma 2.3. In particular, we need only establish the claim that $\mu \circ \Phi \neq \nu_3^1$ and $\mu \circ \Phi \neq \nu_3^2$. For, if $\mu \circ \Phi$ is extraordinary, then by Lemma 5.1 there are $w \in W$ and $i \in \{1, 2\}$ such that $\mu \circ \Phi = \text{Conj}_w \circ \nu_3^i$; taking $g \in A$ with $\mu(g^{-1}) = w$, we find $\mu \circ (\text{Conj}_g \circ \Phi) = \text{Conj}_{w^{-1}} \circ \mu \circ \Phi = \nu_3^i$, so we can apply the claim with $\text{Conj}_g \circ \Phi$ instead of Φ to derive a contradiction.

In order to settle the claim, suppose that $\mu \circ \Phi = \nu_3^1$. Let

$$\begin{aligned} r_1 &= s_1, & r_2 &= s_2, & r_3 &= s_3, \\ r_4 &= s_1 s_2 s_1, & r_5 &= s_2 s_3 s_2, & r_6 &= s_2 s_1 s_2, \\ r_7 &= s_1 s_2 s_3 s_2 s_1, & r_8 &= s_3 s_2 s_1 s_2 s_3, & r_9 &= s_1 s_2 s_1 s_2 s_1, \\ r_{10} &= s_1 s_3 s_2 s_1 s_2 s_3 s_1, & r_{11} &= s_2 s_1 s_2 s_3 s_2 s_1 s_2, & r_{12} &= s_2 s_1 s_3 s_2 s_1 s_2 s_3 s_1 s_2, \\ r_{13} &= s_1 s_2 s_1 s_2 s_3 s_2 s_1 s_2 s_1, & r_{14} &= s_1 s_2 s_1 s_3 s_2 s_1 s_2 s_3 s_1 s_2 s_1, & r_{15} &= s_2 s_1 s_2 s_1 s_3 s_2 s_1 s_2 s_3 s_1 s_2 s_1 s_2. \end{aligned}$$

It is easily verified that $T = \{wsw^{-1}; w \in W, s \in S\} = \{r_1, \dots, r_{15}\}$. Let

$$\tau_2 = \sigma_2 \sigma_1 \sigma_2 \sigma_3 \sigma_2, \quad \tau_3 = \sigma_3 \sigma_2 \sigma_1.$$

A direct calculation gives

$$\begin{aligned} U(\tau_2)(0, r_1) &= (0, r_{12}), & U(\tau_2)(0, r_2) &= (1, r_5), & U(\tau_2)(0, r_3) &= (1, r_9), \\ U(\tau_2)(0, r_4) &= (0, r_{15}), & U(\tau_2)(0, r_5) &= (1, r_{11}), & U(\tau_2)(0, r_6) &= (0, r_4), \\ U(\tau_2)(0, r_7) &= (0, r_8), & U(\tau_2)(0, r_8) &= (1, r_6), & U(\tau_2)(0, r_9) &= (0, r_{13}), \\ U(\tau_2)(0, r_{10}) &= (0, r_3), & U(\tau_2)(0, r_{11}) &= (0, r_1), & U(\tau_2)(0, r_{12}) &= (1, r_2), \\ U(\tau_2)(0, r_{13}) &= (0, r_{14}), & U(\tau_2)(0, r_{14}) &= (0, r_{10}), & U(\tau_2)(0, r_{15}) &= (0, r_7). \end{aligned}$$

and

$$\begin{aligned} U(\tau_3)(0, r_1) &= (1, r_8), & U(\tau_3)(0, r_2) &= (0, r_{10}), & U(\tau_3)(0, r_3) &= (0, r_2), \\ U(\tau_3)(0, r_4) &= (1, r_5), & U(\tau_3)(0, r_5) &= (0, r_{12}), & U(\tau_3)(0, r_6) &= (0, r_7), \\ U(\tau_3)(0, r_7) &= (1, r_3), & U(\tau_3)(0, r_8) &= (0, r_{11}), & U(\tau_3)(0, r_9) &= (0, r_1), \\ U(\tau_3)(0, r_{10}) &= (0, r_6), & U(\tau_3)(0, r_{11}) &= (0, r_{14}), & U(\tau_3)(0, r_{12}) &= (0, r_{15}), \\ U(\tau_3)(0, r_{13}) &= (0, r_4), & U(\tau_3)(0, r_{14}) &= (0, r_9), & U(\tau_3)(0, r_{15}) &= (0, r_{13}). \end{aligned}$$

Since $\mu(\Phi(\sigma_2)) = \nu_3^1(\sigma_2) = s_2 s_1 s_2 s_3 s_2 = \mu(\tau_2)$, and $\mu(\Phi(\sigma_3)) = \nu_3^2(\sigma_3) = s_3 s_2 s_1 = \mu(\tau_3)$, by Corollary 3.8, there exist $b_1, \dots, b_{15}, c_1, \dots, c_{15} \in \mathbf{Z}$ such that

- if $U(\tau_2)(0, r_i) = (k_i, r_j)$, then $U(\Phi(\sigma_2))(0, r_i) = (2b_i + k_i, r_j)$;
- if $U(\tau_3)(0, r_i) = (k'_i, r'_j)$, then $U(\Phi(\sigma_3))(0, r_i) = (2c_i + k'_i, r'_j)$.

Now, the equalities $U(\Phi(\sigma_2 \sigma_3 \sigma_2))(0, r_i) = U(\Phi(\sigma_3 \sigma_2 \sigma_3))(0, r_i)$ for $i = 2, 5, 6, 8, 10, 11$, are equivalent to the equations

$$\begin{aligned} b_2 - b_{10} + b_{12} - c_2 - c_3 + c_5 &= -1, \\ b_5 - b_{12} + b_{14} - c_2 - c_5 + c_{11} &= 0, \\ b_5 + b_6 - b_7 + c_4 - c_6 - c_8 &= -1, \\ b_7 + b_8 - b_{11} - c_1 + c_6 - c_8 &= 0, \\ b_2 - b_6 + b_{10} + c_3 - c_4 - c_{10} &= 0, \\ b_8 + b_{11} - b_{14} + c_1 - c_{10} - c_{11} &= -1. \end{aligned}$$

The sum of these six equations gives $2(b_2 + b_5 + b_8 - c_2 - c_8 - c_{10}) = -3$ which clearly cannot hold. So, $\mu \circ \Phi \neq \nu_3^1$.

Suppose $\mu \circ \Phi = \nu_3^2$. Using the same procedure as before, we find that the equalities $U(\Phi(\sigma_2\sigma_3\sigma_2))(0, r_i) = U(\Phi(\sigma_3\sigma_2\sigma_3))(0, r_i)$ for $i = 2, 4, 7, 8, 10, 12$, are equivalent to the equations

$$\begin{aligned} b_2 - b_7 + b_{15} - c_2 + c_4 - c_{13} &= 0, \\ b_2 + b_4 - b_{15} - c_4 + c_6 - c_{12} &= 0, \\ b_7 - b_{10} + b_{12} - c_7 + c_{13} - c_{14} &= 0, \\ b_3 + b_8 - b_9 - c_2 - c_8 + c_{10} &= -1, \\ -b_3 + b_8 + b_{10} - c_7 - c_{10} + c_{14} &= 0, \\ -b_4 + b_9 + b_{12} - c_6 + c_8 - c_{12} &= 0. \end{aligned}$$

The sum of these six equations gives $2(b_2 + b_8 + b_{12} - c_2 - c_7 - c_{12}) = -1$, a contradiction. So, $\mu \circ \Phi \neq \nu_3^2$. $\square$

Now, we study the remaining cases.

Proposition 5.4. *For each of the Coxeter graphs H_4 , F_4 , E_6 , E_7 , E_8 , all epimorphisms $A \rightarrow W$ are ordinary.*

Proof. Suppose $\varphi : A \rightarrow W$ is a morphism of groups. In a case-by-case analysis, we find all possible morphisms by listing all possible $(\varphi(\sigma_1), \dots, \varphi(\sigma_n))$ up to conjugacy in W . This is done by computing all sequences $(x_1, \dots, x_n) \in W^n$ satisfying $[x_i, x_j]^{m_{i,j}} = [x_j, x_i]^{m_{i,j}}$ for all $i, j \in \{1, \dots, n\}$, up to conjugacy in W , and next verifying whether the x_i generate W . Excluding the ordinary and trivial cases, we can impose some a priori conditions on the x_i . For instance, in view of Lemma 2.1, we may assume that some x_i has order at least 3, and surjectivity of φ implies that some x_i has sign -1 .

It is well known (and easy to verify) that x_i and x_j are conjugate whenever i and j are joined by a path all of whose edges have odd labels. Since F_4 is the only diagram with two connected components with respect to this kind of connectivity, we deal with it first.

F_4 . In view of Lemma 2.1 and symmetry of the diagram, we may assume that x_1 has order greater than 2. We describe a calculational proof using GAP. The group W has 25 conjugacy classes, of which 17 contain elements x_1 of order at least 3. We first compute a complete set of representatives of the conjugacy classes of pairs (x_1, x_2) of elements of W such that x_1 and x_2 are of order at least 3, conjugate and satisfy $x_1x_2x_1 = x_2x_1x_2$. (For each of the 17 representatives x_1 , we compute the orbits of $C_W(x_1)$ in the conjugacy class of x_1 of elements x_2 with $x_1x_2x_1 = x_2x_1x_2$, and take representatives.) There are 53 such representatives. Since none of these generates W , we may assume that x_3 and x_4 are nontrivial. For each pair (x_1, x_2) we compute representatives of the $C_W(x_1, x_2)$ -orbits under conjugation on $C_W(x_1)$ with $[x_2, x_3]^4 = [x_3, x_2]^4$. There are 513 such triples (with $x_3 \neq 1$). Finally, for each such triple, we compute representatives x_4 of each $C_W(x_1, x_2, x_3)$ -orbit of elements conjugate to x_3 , contained in $C_W(x_1, x_2)$, and satisfying $x_3x_4x_3 = x_4x_3x_4$. There are 441 such quadruples (x_1, x_2, x_3, x_4) . Each of these generates a proper subgroup of W . In particular, for F_4 , there are no extraordinary epimorphisms.

For the remaining cases, all x_i will be conjugate and hence have sign -1 . Moreover, from the diagrams it will be clear that we may assume x to be injective when viewed as a map on $\{1, \dots, n\}$ (observe that if i and j are nodes with $x_i = x_j$, then $x_k = x_i$ for each node adjacent to i but not to j [by an odd-labelled edge], so by induction, x would be a constant map, and so the corresponding φ would have a cyclic image, which does not occur). By the same argument, we may even assume x to be strong injective, by which we mean that $\langle x_i \rangle \neq \langle x_j \rangle$ for $i \neq j$.

For $J \subseteq \{1, \dots, n\}$, denote by X_J the set of all W -conjugacy classes of strong injective maps $x : J \rightarrow W$, denoted by $j \mapsto x_j$, such that the x_j ($j \in J$) are all from the same conjugacy class in W , have order at least 3 and satisfy $[x_i, x_j]^{m_{i,j}} = [x_j, x_i]^{m_{i,j}}$ and $\text{sg}(x_i) = -1$ for all $i, j \in J$. The sets X_J can be calculated in GAP by similar computations to those described for F_4 above. Since $X_J = \emptyset$ for some $J \subseteq \{1, \dots, n\}$, there are no extraordinary epimorphisms. We give some of the statistics involved.

H_4 . There are 34 conjugacy classes, but $|X_{\{4\}}| = 8$. We find $|X_{\{2,4\}}| = 6$, $|X_{\{1,2,4\}}| = 6$, and $|X_{\{1,2,3,4\}}| = 0$.

E_6 . There are 25 conjugacy classes, but $|X_{\{1\}}| = 8$. Now $|X_{\{1,2,6\}}| = 11$ and $|X_{\{1,2,4,6\}}| = 0$.

E_7 . There are 60 conjugacy classes, but $|X_{\{1\}}| = 25$, $|X_{\{2,3,5,7\}}| = 840$, and $|X_{\{1,2,3,5,7\}}| = 0$.

E_8 . There are 112 conjugacy classes, but $|X_{\{1\}}| = 43$, $|X_{\{2,3,5,7\}}| = 17700$, and $|X_{\{2,3,5,7,8\}}| = 0$. $\square$

Corollary 5.5. *For each of the Coxeter graphs H_4 , F_4 , E_6 , E_7 , E_8 , the colored Artin group CA is a characteristic subgroup of A . $\square$*

References

- [Art] E. Artin, *Braids and permutations*, Ann. of Math. **48** (1947), 643–649.
- [Bou] N. Bourbaki, *Groupes et algèbres de Lie, Chapitres 4, 5, et 6*, Hermann, Paris, 1968.
- [BS] E. Brieskorn, K. Saito, *Artin-Gruppen und Coxeter-Gruppen*, Invent. Math. **17** (1972), 245–271.
- [Fra] W.N. Franzsen, *Automorphisms of Coxeter groups*, Ph. D. Thesis, University of Sydney, 2001.
- [Lin] V.Y. Lin, *Artin braids and related groups and spaces*, Algebra. Topology. Geometry, Vol. 17, pp. 159–227, 308, Akad. Nauk SSSR, Vsesoyuz. Inst. Nauchn. i Tekhn. Informatsii, Moscow, 1979.
- [Zin] V.M. Zinde, *Some homomorphisms of the Artin groups of the series B_n and D_n into groups of the same series and into symmetric groups*, Uspehi Mat. Nauk **32** (1977), 189–190.

Arjeh M. Cohen
Department of Mathematics and Computer Science
Eindhoven University of Technology
PO Box 513
5600 MB Eindhoven
THE NETHERLANDS
A.M.Cohen@tue.nl

Luis Paris
Laboratoire de Topologie
Université de Bourgogne
UMR 5584 du CNRS, BP 47870
21078 Dijon cedex
FRANCE
lparis@u-bourgogne.fr